25

The invention claimed is:

1. A method of treating and/or preventing metabolic diseases in a patient for whom metformin therapy is inappropriate due to at least one contraindication against metformin comprising orally administering to the patient a DPP-IV 5 inhibitor wherein the contraindication is selected from the group consisting of:

renal disease, renal impairment or renal dysfunction, unstable or acute congestive heart failure, acute or chronic metabolic acidosis, and hereditary galactose intolerance.

- 2. The method according to claim 1 wherein the patient is ineligible for metformin therapy due to contraindication against metformin.
- 3. The method according to claim 1 wherein the patient is in need of reduced dose metformin therapy due to contraindication against metformin.
- **4**. The method according to claim **1** wherein the metabolic disease is type **2** diabetes mellitus.
- 5. The method according to claim 1 wherein the contraindication is renal disease, renal impairment or renal dysfunction.
- 6. The method according claim 1, wherein the DPP-IV inhibitor is

either, in a first embodiment (embodiment A),

of formula (I)

of formula (II)

of formula (III)

of formula (IV)

$$R1$$
 N
 $R2$
 CN
 $R1$
 $R2$
 $R2$
 $R3$

wherein R1 denotes ([1,5]naphthyridin-2-yl)methyl, (quinazolin-2-yl)methyl, (quinoxalin-6-yl)methyl, (4-methyl-quinazolin-2-yl)methyl, 2-cyano-benzyl, (3-cyano-quinolin-2-yl)methyl, (3-cyano-pyridin-2-yl)methyl, (4-methyl-pyrimidin-2-yl)methyl, or (4,6-dimethyl-pyrimidin-2-yl)methyl and R2 denotes 3-(R)-amino-piperidin-1-yl, (2-amino-2-methyl-propyl)-methylamino or (2-(S)-amino-propyl)-methylamino,

or its pharmaceutically acceptable salt;

or, in a second embodiment (embodiment B),

selected from the group consisting of sitagliptin, vildagliptin, saxagliptin, alogliptin,

(2S)-1-{[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethy-lamino]-acetyl}-pyrrolidine-2-carbonitrile,

(2S)-1-{[1,1,-Dimethyl-3-(4-pyridin-3-yl-imidazol-1-yl)-propylamino]-acetyl}-pyrrolidine-2-carbonitrile,

(S)-1-((2S,3S,11bS)-2-Amino-9,10-dimethoxy-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-3-yl)-4-fluoromethyl-pyrrolidin-2-one,

(3,3-Difluoropyrrolidin-1-yl)-((2S,4S)-4-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrrolidin-2-yl)methanone,

(1((3S,4S)-4-Amino-1-(4-(3,3-difluoropyrrolidin-1-yl)-1,3,5-triazin-2-yl)pyrrolidin-3-yl)-5,5-difluoropiperidin-2-one,

(2S,4S)-1-{2-[(3S,1R)-3-(1H-1,2,4-Triazol-1-ylmethyl)cyclopentylamino]-acetyl}-4-fluoropyrrolidine-2-carbonitrile.

(R)-2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4fluoro-benzonitrile,

5-{(S)-2-[2-((S)-2-Cyano-pyrrolidin-1-yl)-2-oxo-ethy-lamino]-propyl}-5-(1H-tetrazol-5-yl)-10,11-dihy-dro-5H-dibenzo[a,d]cycloheptene-2,8-dicarboxylic acid bis-dimethylamide,

3-{(2S,4S)-4-[4-(3-Methyl-1-phenyl-1H-pyrazol-5-yl) piperazin-1-yl]pyrrolidin-2-ylcarbonyl}thiazolidine,

[(2R)-1-{[(3R)-Pyrrolidin-3-ylamino] acetyl}pyrrolidin-2-yl]boronic acid,

(2S,4S)-1-[2-[(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile,

2-({6-[(3R)-3-amino-3-methylpiperidin-1-yl]-1,3-dimethyl-2,4-dioxo-1,2,3,4-tetrahydro-5H-pyrrolo[3,2-d]pyrimidin-5-yl}methyl)-4-fluorobenzonitrile, and

6-[(3R)-3-amino-piperidin-1-yl]-5-(2-chloro-5-fluoro-benzyl)-1,3-dimethyl-1,5-dihydro-pyrrolo[3,2-d]pyrimidine-2,4-dione,

or its pharmaceutically acceptable salt.

7. The method according claim 1, wherein said DPP-4 inhibitor is selected from the group consisting of

1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-(3-(R)-amino-piperidin-1-yl)-xanthine,

1-[([1,5]naphthyridin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

1-[(quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

2-((R)-3-amino-piperidin-1-yl)-3-(but-2-ynyl)-5-(4-methyl-quinazolin-2-ylmethyl)-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,

1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(2-amino-2-methyl-propyl)-methylamino]-xanthine,

1-[(3-cyano-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

1-(2-cyano-benzyl)-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,